

RESEARCH ARTICLE

# Amphibian skin-associated *Pigmentiphaga*: Genome sequence and occurrence across geography and hosts

Molly C. Bletz<sup>1,2\*</sup>, Boyke Bunk<sup>3</sup>, Cathrin Spröer<sup>3</sup>, Peter Biwer<sup>4</sup>, Silke Reiter<sup>5</sup>, Falitiana C. E. Rabemananjara<sup>6</sup>, Stefan Schulz<sup>4</sup>, Jörg Overmann<sup>3,7</sup>, Miguel Vences<sup>2</sup>

**1** Department of Biology, University of Massachusetts Boston, Boston, MA, United States of America, **2** Zoological Institute, Technische Universität Braunschweig, Braunschweig, Germany, **3** DSMZ, German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany, **4** Institute of Organic Chemistry, Technische Universität Braunschweig, Braunschweig, Germany, **5** Institute for Insect Biotechnology, Justus Liebig University Giessen, Giessen, Germany, **6** Department of Zoology and Animal Biodiversity, University of Antananarivo, Antananarivo, Madagascar, **7** Microbiology Institute, Technische Universität Braunschweig, Braunschweig, Germany

\* [molly.bletz@gmail.com](mailto:molly.bletz@gmail.com)



## OPEN ACCESS

**Citation:** Bletz MC, Bunk B, Spröer C, Biwer P, Reiter S, Rabemananjara FCE, et al. (2019) Amphibian skin-associated *Pigmentiphaga*: Genome sequence and occurrence across geography and hosts. PLoS ONE 14(10): e0223747. <https://doi.org/10.1371/journal.pone.0223747>

**Editor:** Bi-Song Yue, Sichuan University, CHINA

**Received:** May 17, 2019

**Accepted:** September 29, 2019

**Published:** October 11, 2019

**Copyright:** © 2019 Bletz et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** The complete genome sequence of *Pigmentiphaga aceris* Mada1488 has been deposited at NCBI GenBank under the accession no CP043046. (BioProject number PRJNA561098).

**Funding:** The sampling and culturing of amphibian skin bacteria in Madagascar was supported by a grant from the Mohamed bin Zayed Conservation Fund to MB and RH, a grant from the Amphibian Survival Alliance to MB, RH, and MV, a grant from Chester Zoo to MB and RH, a scholarship of the

## Abstract

The bacterial communities colonizing amphibian skin have been intensively studied due to their interactions with pathogenic chytrid fungi that are causing drastic amphibian population declines. Bacteria of the family *Alcaligenaceae*, and more specifically of the genus *Pigmentiphaga*, have been found to be associated specifically to arboreal frogs. Here we analyze their occurrence in a previously assembled global skin microbiome dataset from 205 amphibian species. *Pigmentiphaga* made up about 5% of the total number of reads in this global dataset. They were mostly found in unrelated arboreal frogs from Madagascar (Mantellidae and Hyperoliidae), but also occurred at low abundances on Neotropical frogs. Based on their 16S sequences, most of the sequences belong to a clade within *Pigmentiphaga* not assignable to any type strains of the five described species of the genus. One isolate from Madagascar clustered with *Pigmentiphaga aceris* (>99% sequence similarity on 16S rRNA gene level). Here, we report the full genome sequence of this bacterium which, based on 16S sequences of >97% similarity, has previously been found on human skin, floral nectar, tree sap, stream sediment and soil. Its genome consists of a single circular chromosome with 6,165,255 bp, 5,300 predicted coding sequences, 57 tRNA genes, and three rRNA operons. In comparison with other known *Pigmentiphaga* genomes it encodes a higher number of genes associated with environmental information processing and cellular processes. Furthermore, it has a biosynthetic gene cluster for a nonribosomal peptide synthetase, and bacteriocin biosynthetic genes can be found, but clusters for  $\beta$ -lactones present in other comparative *Pigmentiphaga* genomes are lacking.

German Academic Exchange Service (DAAD) to MB, and a grant from the Deutsche Forschungsgemeinschaft (DFG) to MV (VE247/9-1). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing interests:** The authors have declared that no competing interests exist.

## Introduction

The cutaneous microbiome of amphibians has become a well-studied system, triggered by the rise of the pathogenic fungi, *Batrachochytrium dendrobatidis* (*Bd*) and *B. salamandrivorans* (*Bsal*). These fungi colonize the amphibian skin and are causing drastic population declines and extinctions in this class of animals [1,2]. The bacterial communities associated to amphibians interact with these fungi and some of these bacteria have the potential to inhibit the growth of *Bd* and *Bsal*, thus providing protection to their hosts [3].

Recent research based on next-generation amplicon sequencing of the 16S rRNA gene show that bacterial communities on the skin of amphibians are predominantly composed of common bacteria recruited from environmental reservoirs [4], and their dominant members can be readily cultured [5]. Unsurprisingly these communities are strongly controlled by environmental factors, e.g., bioclimate [6] and microhabitat [7–9]. However, clear differences have also been found between co-occurring hosts [7,10,11], suggesting that the skin mucosal differences among amphibian species act as filters determining which bacterial species are recruited into the community.

Considering the strong environmental influences on the amphibian cutaneous microbiome, it is of particular interest to analyze in more depth those bacteria that regularly colonize this habitat but are restricted to certain host taxa or host ecomorphs. An in-depth understanding of the genomic background, variation, phylogenetic relationships, and distribution of these bacteria may offer clues to understand which traits predispose them to successfully colonize this particular habitat.

This study was triggered by the observation that operational taxonomic units (OTUs) of the family *Alcaligenaceae* were strongly associated to arboreal ecomorphs in a study on Madagascan amphibians [12], and were also found to be a common member of the cutaneous microbiome of several Central American tree frogs, such as *Agalychnis callidryas* [13,14]. The family was represented by a pure culture isolate identified as *Pigmentiphaga* by 16S sequences in our bacterial culture collection from Madagascar frog skin [15], and also the sequences of *Alcaligenaceae* OTUs identified by amplicon sequences from Madagascar frogs.

*Pigmentiphaga* is a genus of the family *Alcaligenaceae*, assigned to the order Burkholderiales order within the Betaproteobacteria, and currently containing five species: the type species, *P. kullae*, plus *P. aceris*, *P. daeguensis*, *P. litoralis*, and *P. solis* [16–20]. According to these descriptions, the genus contains gram-negative, facultatively anaerobic, motile or nonmotile, catalase- and oxidase-positive, rod-shaped bacteria, found in diverse environments: *P. daeguensis* from dye wastewater, *P. litoralis* from tidal sediment, *P. soli* from soil, *P. aceris* from tree sap [17–20], and an unidentified species from tree-associated nematodes [21]. Furthermore, *Pigmentiphaga* have also been isolated from human clinical material [22], and genome sequences are available from these isolates [23]. *Pigmentiphaga* have been studied in the context of their ability to degrade azo dyes and aniline [16,24], and their role also has been discussed in the context of biphenyl-degradation [25].

Here, we analyze the occurrence of OTUs assigned to *Pigmentiphaga* on the skin of amphibians across taxa, geography and ecomorphs, assemble the full genome sequence of one *Pigmentiphaga* isolate obtained from a Madagascan frog, and analyze the phylogenetic relationships and differentiation of this isolate.

## Methods

### Analysis of amplicon data

To explore the distribution of the focal bacterial groups on amphibian hosts we used a recently published global dataset of amphibian skin microbiomes [6], and extracted all OTUs assigned

to the family *Alcaligenaceae* and genus *Pigmentiphaga*. This global dataset is a compilation of targeted amplicon sequencing data of the 16S rRNA gene (V4 region) generated on Illumina sequencing platforms. It contains skin microbiome samples from 2,349 post-metamorphic amphibians, comprising 27 amphibian families (205 species) collected from 12 countries (5 continents) [6]. In this study, quality-filtered Illumina reads were classified into sub-operational taxonomic units (sOTUs) using the deblur pipeline [26]. These data were subsequently rarified and taxonomy was assigned using the Ribosomal Database Classifier [27] in QIIME [28] (see [6] for detailed methods). We examined the distribution of the genus *Pigmentiphaga* across locations (i.e. countries), host species, and host ecomorphological classes. To explore the distribution of the focal bacteria across host phylogeny, a phylogenetic tree of arboreal and scansorial host genera was built using timetree.org [29]. All plots were produced with ggplot2 [30].

### Isolate sampling

The bacterium for which we here report the full genome sequence was isolated on 1% tryptone agar from a skin swab of a single individual of the amphibian species, *Mantella crocea* at Torotorofotsy "Prolemur Camp" near Andasibe, Madagascar (-18.7709 S, 48.43222 E). This individual frog was captured by gloved hands and kept in a sterile Whirl-Pak bag for no longer than 1 hour before sampling. Frog skin microbiota were sampled by first rinsing the skin to remove transient microbes and then swabbing the ventral skin with a sterile rayon tipped swab 10 times. Swabs were stored in Tryptic Soy Yeast Extract media with 20% Glycerol and frozen until processing. The frog was immediately released at the site of capture after sampling.

### DNA extraction and complete genome sequencing

DNA was isolated using Qiagen Genomic-tip 100/G (Qiagen, Hilden Germany) according to manufacturer instructions. SMRTbell™ template library was prepared according to the instructions from PacificBiosciences, Menlo Park, CA, USA, following the Procedure & Checklist—Greater Than 10 kb Template Preparation. Briefly, for preparation of 15kb libraries 8μg genomic DNA was sheared using g-tubes™ from Covaris, Woburn, MA, USA according to the instructions of the manufacturer. DNA was end-repaired and ligated overnight to hairpin adapters applying components from the DNA/Polymerase Binding Kit P6 from Pacific Biosciences, Menlo Park, CA, USA. Reactions were carried out according to the manufacturer instructions. BluePippin™ Size-Selection to greater than 4 kb was performed according to the manufacturer's instructions (Sage Science, Beverly, MA, USA). Conditions for annealing of sequencing primers and binding of polymerase to purified SMRTbell™ template were assessed with the Calculator in RS Remote, PacificBiosciences, Menlo Park, CA, USA. 1 SMRT cell was sequenced on the PacBio RSII (PacificBiosciences, Menlo Park, CA, USA) taking one 240-minutes movie. SMRT sequencing revealed a total number of 85,590 reads with a mean read length of 12,138 bp and a N50 value of 16,449 bp. From the same batches of DNA, short insert libraries were created using the Illumina Nextera XT DNA Library Prep Kit (Illumina, San Diego, CA, USA) and sequenced on an Illumina MiSeq (Illumina, San Diego, CA, USA) resulting in 7,916,070 paired-end reads of 2x76 bp.

### Genome assembly and annotation

Genome assembly was performed applying the RS\_HGAP\_Assembly.3 protocol included in SMRT Portal version 2.3.0 using default parameters. The assembly revealed a single circular chromosome with a coverage of 133x. The chromosome was circularized, artificial redundancies at the ends of the contigs were removed and adjusted to *dnaA* as the first gene. Error-correction was performed by a mapping of Illumina short reads onto finished genome using

Burrows-Wheeler Alignment bwa 0.6.2 in paired-end (sample) mode using default setting [31] with subsequent variant and consensus calling using VarScan 2.3.6 (Parameters: mpileup2cns —min-coverage 10—min-reads2 6—min-avg-qual 20—min-var-freq 0.8—min-freq-for-hom 0.75—p-value 0.01—strand-filter 1—variants 1—output-vcf 1) [32]. A consensus concordance of QV60 could be reached. Automated genome annotation was carried out using Prokka [33] and NCBI PGAP [34].

## Genome comparisons

We used Mauve software [35] to compare gene arrangement of the newly sequenced genome with those of other available *Pigmentiphaga* genomes. We compared the general gene functional characterization (KEGG functional categories) of the newly sequenced genomes with all other available *Pigmentiphaga* genomes using BlastKOALA [36]. On average, 50% of genomes' protein coding sequences were annotated. Identification of natural product gene clusters was performed with the antibiotics and Secondary Metabolite Analysis SHell (antiSMASH version beta5; <https://antismash.secondarymetabolites.org>) [37]. AntiSMASH is an online platform that allows for a genome-wide identification and analysis of secondary metabolite BGCs in bacterial genomes, by integrating and cross-linking with a large number of *in silico* secondary metabolite analysis tools like CLUSEAN [38], BAGEL2 [39], ClustScan [40], and NORINE [41].

## Phylogenetic analysis

We performed BLAST searches against the NCBI database, using the full 16S rRNA gene sequence from the sequenced genome to understand the distribution of this bacterium outside of amphibian hosts. We also used the MOLE-BLAST tool of NCBI to retrieve from GenBank the sequences of bacterial taxa most closely related to the *Alcaligenaceae* sOTUs in our amplicon data set. MOLE-BLAST is an experimental tool to find closest database neighbors of submitted query sequences, by computing a multiple sequence alignment (MSA) between the query sequences along with their top BLAST database hits.

The obtained sequences, along with sequences of our isolates and amplicon-derived sOTU sequences, were aligned with the MAFFT 7.0 algorithm [42] and phylogenetically analyzed in MEGA v. 7 [43]. We successively filtered the data set to remove highly deviant sequences retrieved from the database (as well as sOTU sequences associated to them), as identified by obvious alignment artefacts and excessively long branches in exploratory phylogenetic trees. The final Maximum Likelihood (ML) tree was computed under the GTR+G model of sequence evolution as determined under the Bayesian Information criterion in MEGA 7.

## Volatile compound analysis

A bacterial culture of the target bacterial strain was incubated on 1% tryptone agar for seven days at room temperature. Headspace extracts were obtained using a vacuum pump to draw clean air (purification by active charcoal filter) through a 250 mL glass vessel containing the culture plate. The air was then passed through a thermal desorption tube filled with an absorbent (Tenax TA Tube; GERSTEL, Mülheim an der Ruhr, Germany) for 5 h (three replicates).

Thermal desorption tubes were desorbed using a thermal desorption unit (TDU), cooled injection system (CIS) and a MultiPurposeSampler (MPS) autosampler (GERSTEL, Mülheim an der Ruhr, Germany) connected to an Agilent 7890B gas chromatograph. The gas chromatograph was equipped with a HP-5 MS fused silica capillary column (30 m, 0.25 i. d., 0.25  $\mu$ m film, Hewlett-Packard, Wilmington, USA) connected to an Agilent 5977A mass-selective detector. Conditions: transfer line 300°C, electron energy 70 eV. Thermal desorption: 30°C, increasing at 60°C/min to 280°C (10 min isothermal). Cooled injection: -150°C, increasing at

12°C/min to 300°C (3 min isothermal). Gas chromatographic method: 50°C (5 min isothermal), increasing at 5°C/min to 320°C, and operated in splitless mode. Helium was used as carrier gas at 1.2 ml/min. GC retention indices (RI) were determined from a homologous series of *n*-alkanes (C<sub>8</sub>–C<sub>30</sub>). Compounds were identified by comparison of mass spectra and retention indices with those of authentic samples.

## Results and discussion

### Representation of *Alcaligenaceae* in amphibian cutaneous microbiomes

In the global dataset of amplicon sequences from 205 amphibian species [6] *Alcaligenaceae* sOTUs made up 284,771 out of 5,872,500 total rarified reads in the final data set (4.8%). *Alcaligenaceae* sOTUs (at a minimum threshold of 5 reads) were found in a total of 119 amphibian species from eight countries, and *Pigmentiphaga* sOTUs in 95 amphibian species. In a culture database of amphibian skin bacteria [22, Bletz & Woodhams unpublished data] 28 of 5938 isolates were from the *Alcaligenaceae*, and our isolate was the sole member from the genus *Pigmentiphaga*. Therefore, apparently, *Alcaligenaceae* and more specifically *Pigmentiphaga* appear to be underrepresented in culture databases of amphibian skin microbiota and thus might be less readily culturable than other bacteria from this habitat [5].


The family *Alcaligenaceae* currently contains 27 genera (UniProt 2019); in the amphibian microbiome data set, 338 out of a total of 124,348 sOTUs were assigned to this family, and of these, 35 to the genus *Pigmentiphaga* (reads = 266,723). The remaining *Alcaligenaceae* reads were assigned to the genera *Achromobacter* (n = 3,355), *Alcaligenes* (n = 666), *Sutterella* (n = 20) *Oligella* (n = 16), *Denitrobacter* (n = 3), *Candidimonas* (n = 12) or were left unassigned to a specific genus (n = 13,976).

Confirming previous findings [12–14], in our global skin microbiome dataset [6] *Pigmentiphaga* was predominantly found on arboreal species as well as scansorial hosts within the amphibian clades Mantellidae (mean: 7.7% + 1.5%SD) and Hyperoliidae (mean: 7.9% + 4.7% SD, which are distributed in Madagascar, and (Hyperoliidae only) in mainland Africa (Fig 1). The genus also appears on amphibians from the genus *Pseudacris* (Fig 1). Overall, *Pigmentiphaga* was more common on amphibians from Madagascar; however, this could be associated with extensive sampling of arboreal hosts within this country.

### Phylogenetic diversity of *Pigmentiphaga* in amphibian cutaneous microbiomes

We aligned short amplicon-based consensus sequences of *Alcaligenaceae* amplicons with the longer 16S sequences of all Madagascan amphibian-derived isolates from a previous work [15] belonging to this family. We then used a series of BLAST and MOLE-BLAST searches, allowing for hits with and without environmental sequences, and restricting searches to type strains or not, to retrieve a representative set of 273 homologous *Alcaligenaceae* sequences of the 16S rRNA gene for analysis of phylogeny and environmental distribution of our focal bacterial taxa. The exploratory analysis of these sequences along with our *Alcaligenaceae* sOTU and isolate sequences placed 92 sOTUs, 8 isolates and 82 related sequences retrieved from GenBank in a clade with the *Pigmentiphaga* type strains. A ML tree calculated on this restricted dataset (Fig 2) reveals a large diversity of *Pigmentiphaga*, many of which are not assignable to any of the described species. A large number of 61 additional sOTUs from the amphibian skin, as well as one isolate from the skin of a fire salamander (DE946; accession number MH512662) and two from Madagascar frogs (Mada281, Mada1835; accession numbers MF526411, MF523827), are placed in a large subclade of putative *Pigmentiphaga* sequences that does not contain any type






**Fig 1. Distribution of *Pigmentiphaga* spp. across amphibian hosts.** relative abundance within amphibian skin microbiomes across host eco-morphology classes (A), countries (B), and phylogeny of arboreal and scansorial amphibian hosts (C).

<https://doi.org/10.1371/journal.pone.0223747.g001>

strain sequences. This subclade also contains sOTU7503, the most widespread *Alcaligenaceae* sOTU in our global amplicon-derived data set (45,697 reads). Various sequences retrieved from GenBank and included in this subclade are named *P. daguensis* but are unlikely to belong to this species, given that the type strain is placed in another, phylogenetically distant clade. Whether this diverse subclade is to be assigned to *Pigmentiphaga* definitively, or to another, possibly undescribed genus in the *Alcaligenaceae*, will require additional study.

### Genome characteristics of *Pigmentiphaga aceris* isolated from amphibian skin

One of our isolates (Mada1488) was placed close to *P. aceris* and had >99% sequence similarity with the type strain of this species (Table 1). The sequence obtained by direct Sanger



**Fig 2. Maximum likelihood phylogenetic tree of selected *Pigmentiphaga* based on DNA sequences of up to 1478 bp of the bacterial 16S rRNA gene.** Sequences of amphibian skin bacteria from an Illumina-based amplicon survey<sup>[6]</sup> are colored purple; isolates from amphibian skin are colored blue. Red color highlights the sequences of the *P. aceris* strain used for genome sequencing. Sequences from type or reference strains are boldfaced. *Alcaligenes faecalis*, the type species of the type genus of *Alcaligenaceae*, was used as the outgroup. Results of a bootstrap analysis (100 replicates) are marked by gray (bootstrap proportion >50%) and black circles (>70%). Note that due to the inclusion of many short sequences from Illumina amplicon analysis, most nodes did not receive strong bootstrap support.


<https://doi.org/10.1371/journal.pone.0223747.g002>

Table 1. NCBI database matches (97–100% sequence identity) to the 16S rRNA gene of *Pigmentiphaga aceris* (strain Mada1488).

Accession #	Description	Query cover	Percent match	Isolation Source
HM276147.1	Uncultured bacterium clone ncd518h02c1	89%	99.4%	Human skin
NR_157990.1	<i>Pigmentiphaga aceris</i> strain SAP-32	94%	99.3%	tree sap
HM270483.1	Uncultured bacterium clone ncd266g01c1	89%	99.2%	human skin
FJ849553.1	Uncultured bacterium clone SedUMA34	95%	99.2%	arctic stream sediment; ultramafic lithology
JX067670.1	Alcaligenaceae bacterium SAP706.3	98%	99.1%	floral nectar
FN421876.1	Uncultured bacterium, clone 2_F01	92%	99.1%	phyllosphere of clover
HM316492.1	Uncultured bacterium clone ncd307b11c1	89%	99.0%	human skin
JX067692.1	Alcaligenaceae bacterium SAP773.2	98%	99.0%	floral nectar
HM316454.1	Uncultured bacterium clone ncd306e07c1	89%	99.0%	human skin
FJ849534.1	Uncultured bacterium clone SedUMA17 1	95%	98.4%	arctic stream sediment; ultramafic lithology
MH667611.1	<i>Pigmentiphaga</i> sp. strain IMT-318	94%	97.1%	soil (USA)

<https://doi.org/10.1371/journal.pone.0223747.t001>

sequencing of DNA extracted from this isolate agreed fully with the sequence from both 16S copies found in the assembled genome. One amplicon-derived sOTU also the sequence of this isolate. In our global amphibian data set, 139 of these reads came from the sOTU matching the Mada1488 isolate. This sOTU was found on frogs of the genera *Anaxyrus*, *Boophis*, *Colostethus*, *Craugastor*, *Gephyromantis*, *Eleutherodactylus*, *Lithobates*, *Mantidactylus*, *Mantella*, and *Plethodontohyla*. Thus, the bacterium represented by our culture (Mada1488) was not



**Fig 3. The circular genome of 6,165,255 bp of *Pigmentiphaga aceris* (strain Mada1488).** In blue (circle 1) genes lying on the forward strand are shown and in red (circle 2) those on the reverse strand. In circle 3 tRNA genes are shown in brown, often clustered together with green rRNA genes. Circle 4 shows the GC content  $(G+C)/(A+T+G+C)$ , whereas in circle 5 a GC skew  $(G-C)/(G+C)$  is shown. This map has been created using DNAPlotter [44].

<https://doi.org/10.1371/journal.pone.0223747.g003>



very common in our Illumina dataset. As it was the only cultured *Pigmentiphaga* from amphibian skin assignable to a known species we nevertheless chose this isolate for genome sequencing, to obtain first data of the genomic background of these bacteria.

In culture, this bacterium forms white, glossy colonies with circular form; the elevation is raised and the margins are entire. In a growth inhibition assay (see [15] for methods) the metabolites produced by this bacterium reduced the growth of the amphibian skin pathogen *Batrachochytrium dendrobatidis* by 60%. BLAST searches in the NCBI nucleotide database revealed that bacterial strains with >97% 16S sequence identity to Mada1488 have been found on human skin, in floral nectar, tree sap, arctic stream sediment, and soil (Table 1). The highest identity was found with an uncultured isolate from human skin (99.4% identity), directly followed by the *P. aceris* type strain (SAP-32) with 99.3% identity.

The complete genome of *Pigmentiphaga aceris* (Mada1488) consists of a single circular chromosome with 6,165,255 bp and a GC content of 62.1%. Prokka predicted 5,300 coding sequences, 57 tRNA genes, and three rRNA operons (Fig 3). A multiple genome alignment suggests that the new genome shows only limited similarities to the five congeneric genomes available (S1 Fig); however, the five available genomes all belong to closely related strains (all >98% 16S similarity to the type strains of *P. kullae* and *P. daeguensis*, which themselves show 99.6% similarity, questioning the distinctness of these two taxa at the species level).

The newly sequenced genome of *P. aceris* (Mada1488) differs in its gene functions compared to other known *Pigmentiphaga* genomes; more specifically, genes associated with environmental information processing and cellular processes were more abundant in Mada1488 (S2 Fig). Natural product biosynthetic gene clusters (BGCs) are moderately represented in the newly sequenced genome, and the cluster content differs from that in the other available *Pigmentiphaga* genomes (S1 Table). Amongst others, *P. aceris* (Mada1488) contains a nonribosomal peptide synthetase (NRPS) cluster most likely involved in the production of a peptide siderophore similar to enterobactin [45], and a BGC coding for bacteriocin biosynthetic genes, both missing in the other *Pigmentiphaga*. Bacteriocines are widely occurring, ribosomally produced antimicrobial peptides, presumably with an anti-competitor function [46] and an often narrow activity range against Gram-positive as well as Gram-negative bacteria [47,48]. The most noticeable feature in the comparative *Pigmentiphaga* genomes but absent in *P. aceris* (Mada1488) are clusters for  $\beta$ -lactones, a class of protein-inhibiting natural products with a wide range of activities [49]. Furthermore, *P. aceris* (Mada1488) lacks genes involved in ectoin biosynthesis. Because ectoins are usually produced to protect the bacteria from environmental extremes like hyperosmotic conditions [50], the lack of ectoins might support this strain being adapted to rather stable environmental conditions.

### Volatile compounds produced by *Pigmentiphaga aceris*

The ability to produce volatile organic compounds (VOCs) is widespread among bacteria [51] and our data demonstrate this ability also in *Pigmentiphaga aceris* (Mada1488). GC/MS analysis of headspace extracts from this bacterium revealed the release of 21 compounds (S1 Table), among them sulfur-containing volatiles such as methanethiole, dimethyl disulfide (1), dimethyl trisulfide (2), S-methyl ethanethioate (3), S-methyl propanethioate (4), S-methyl 2-methylpropanethioate (5), S-methyl 3-methylbutanethioate (6) and S-methyl phenylethanethioate (7), as well as  $\gamma$ -decalactone (8) (S3 Fig). Some these compounds, for example, dimethyl trisulfide and S-methyl 3-methylbutanethioate, have been associated with inhibition of a variety of plant pathogens [52,53], suggesting they may play a role in suppression of amphibian fungal pathogens, such as *Bd* and *Bsal*.

## Conclusions

To our knowledge, *Pigmentiphaga aceris* (Mada1488) is the first amphibian skin-derived bacterial isolate with a full genome sequenced. Overall, the genome does not present any outstanding characteristics, in line with the hypothesis that the amphibian cutaneous microbiome mainly consists of generalist species recruited from environmental reservoirs. It is remarkable that bacterial strains most similar to Mada1488 have multiple times been found in plant-associated microbiomes (tree sap, nectar, phyllosphere). The finding of a *Pigmentiphaga* similar to the strain described herein on human skin may be explained by a bias of information in genetic databases towards human-derived microbes, but also confirms that these bacteria are not strictly associated to plants only. Yet, it is tempting to relate the apparent common occurrence of *Pigmentiphaga* on plants to its high abundance in treefrogs which may have acquired it from plant-associated reservoirs. To test this hypothesis, in-depth analysis of additional bacteria differentially abundant on arboreal vs. terrestrial amphibians may be rewarding. A wider sampling of genomes represented in amphibian cutaneous microbiomes will be a crucial step to better understand functional properties of these bacterial communities and their potential role in defense against pathogens.

## Data availability

The complete genome sequence of *Pigmentiphaga aceris* Mada1488 has been deposited at NCBI GenBank under the accession no. CP043046. The version described in this paper is the first version. (BioProject no. PRJNA561098). The 16S sequence of this isolate is archived under accession no. MF525803.1. Accession numbers of sequences used in the phylogenetic analysis are given in the respective tree. Accession numbers for the raw data of amplicon analyses are summarized in a previous study [6].

## Supporting information

**S1 Fig. Comparison of genomic composition of the *Pigmentiphaga aceris* strain isolated from amphibian skin (A) to the five other *Pigmentiphaga* genomes available: (A) *Pigmentiphaga* sp. H8, (B) *P. sp.* NML030171, (C) *P. sp.* NML080357, (D) *P. kullae* K24, (E) *P. sp.* IMT-318.** The figure shows a multiple genome alignment calculated with Mauve (Darling et al. 2004), using A as reference. Colinear blocks are indicated by identical colors and indicate homologous DNA regions shared by two or more genomes without sequence rearrangements, and are indicated below the black horizontal line if representing reverse complements of the respective sequence of the reference. Note similarities between genomes A-C, larger differences of D and E, and massive differences in the arrangement of the newly sequenced *P. aceris* genome (F).  
(JPG)

**S2 Fig. Comparative summary of gene function across the newly sequenced *Pigmentiphaga aceris* (red box) and other available genomes from this genus.** Pie charts were created directly from BlastKOALA. Colors for a given functional categories are consistent across each chart; categories are ordered by abundance within a given pie chart.  
(PDF)

**S3 Fig. Selected volatile compounds released by *Pigmentiphaga aceris* (Mada1488):** methanethiole, dimethyl disulfide (1), dimethyl trisulfide (2), S-methyl ethanethioate (3), S-methyl propanethioate (4), S-methyl 2-methylpropanethioate (5), S-methyl 3-methylbutanethioate (6) and S-methyl phenylethanethioate (7), as well as  $\gamma$ -decalactone (8).  
(PDF)

**S1 Table. Natural product biosynthetic gene clusters (BGCs) in available *Pigmentiphaga* genomes as predicted by AntiSMASH.**

(DOCX)

**S2 Table. Volatile compounds released by *Pigmentiphaga aceris* (Mada1488).** Numbered compounds 1–8 are those shown in [S2 Fig](#).

(DOCX)

## Acknowledgments

We are grateful to the Malagasy authorities for issuing research and export permits for this research. We are indebted to numerous local guides and field assistants that help during field work. We thank Simone Severitt and Carola Berg for technical assistance with laboratory work related to genome sequencing. We thank the Malagasy authorities for permits to collect, export and analyze the amphibian-skin derived bacteria, including full genome sequencing (research authorizations 105N-EA04/MG17 from 25 April 2017; collecting permits 182/13/MEF/SG/DGF/DCB.SA/SCB from 1 August 2013, 248/16/MEEF/SG/DGF/DSAP/SCB.Re from 14 October 2016, and 282/16/MEEF/SG/DGF/DSAP/SCB from 28 November 2016). The work was carried out in the framework of a collaboration accord of the Technische Universität Braunschweig with the Cellule d'Urgence Chytride de Madagascar and the Université de Madagascar (Mention Biodiversité Animale), with a Material Transfer Agreement (002/ZBA/17/ZR) from 3 January 2017.

## Author Contributions

**Conceptualization:** Molly C. Bletz, Boyke Bunk, Cathrin Spröer, Falitiana C. E. Rabemananjara, Stefan Schulz, Jörg Overmann, Miguel Vences.

**Data curation:** Molly C. Bletz, Boyke Bunk, Cathrin Spröer, Silke Reiter, Miguel Vences.

**Formal analysis:** Molly C. Bletz, Boyke Bunk, Peter Biwer, Silke Reiter, Stefan Schulz, Miguel Vences.

**Funding acquisition:** Falitiana C. E. Rabemananjara, Miguel Vences.

**Methodology:** Cathrin Spröer, Peter Biwer, Silke Reiter, Stefan Schulz.

**Project administration:** Jörg Overmann.

**Visualization:** Molly C. Bletz.

**Writing – original draft:** Molly C. Bletz, Miguel Vences.

**Writing – review & editing:** Molly C. Bletz, Boyke Bunk, Peter Biwer, Silke Reiter, Falitiana C. E. Rabemananjara, Stefan Schulz, Jörg Overmann, Miguel Vences.

## References

1. Fisher MC, Garner TW, Walker SF. Global emergence of Batrachochytrium dendrobatidis and amphibian chytridiomycosis in space, time, and host. *Annu Rev Microbiol.* 2009; 63: 291–310. <https://doi.org/10.1146/annurev.micro.091208.073435> PMID: 19575560
2. Stegen G, Pasmans F, Schmidt BR, Rouffaer LO, Van Praet S, Schaub M, et al. Drivers of salamander extirpation mediated by Batrachochytrium salamandrorans. *Nature.* Nature Publishing Group; 2017; 544: 353–356. <https://doi.org/10.1038/nature22059> PMID: 28425998

3. Bletz MC, Loudon AH, Becker MH, Bell SC, Woodhams DC, Minbiole KPC, et al. Mitigating amphibian chytridiomycosis with bioaugmentation: Characteristics of effective probiotics and strategies for their selection and use. *Ecol Lett*. 2013; 16. <https://doi.org/10.1111/ele.12099> PMID: 23452227
4. Loudon AH, Woodhams DC, Parfrey LW, Archer HM, Knight R, McKenzie V, et al. Microbial community dynamics and effect of environmental microbial reservoirs on red-backed salamanders (*Plethodon cinereus*). *ISME J*. Nature Publishing Group; 2014; 8: 830–40. <https://doi.org/10.1038/ismej.2013.200> PMID: 24335825
5. Walke JB, Becker MH, Hughey MC, Swartwout M, Jensen R V, Belden LK. Most of the dominant members of amphibian skin bacterial communities can be readily cultured. *Appl Environ Microbiol*. 2015; <https://doi.org/10.1128/AEM.01486-15> PMID: 26162880
6. Kueneman JG, Bletz MC, Becker CG, Woodhams DC, Vences M. Community richness of amphibian skin bacteria correlates with bioclimate at the global scale. *Nat Ecol Evol*. 2019;
7. Kueneman JG, Parfrey LW, Woodhams DC, Archer HM, Knight R, McKenzie VJ. The amphibian skin-associated microbiome across species, space and life history stages. *Mol Ecol*. 2014; 23: 1238–1250. <https://doi.org/10.1111/mec.12510> PMID: 24171949
8. Bletz MC, Perl RGB, Vences M. Skin microbiota differs drastically between co-occurring frogs and newts. *Open Sci*. 2017; 4: 170107.
9. Jani AJ, Briggs CJ. Host and aquatic environment shape the amphibian skin microbiome but effects on downstream resistance to the pathogen *Batrachochytrium dendrobatidis* are variable. *Front Microbiol*. 2018; 9: 1–17. <https://doi.org/10.3389/fmicb.2018.00001>
10. Sabino-Pinto J, Bletz MC, Islam MM, Shimizu N, Bhuju S, Geffers R, et al. Composition of the cutaneous bacterial community in Japanese amphibians: effects of captivity, host species, and body region. *Microb Ecol. Microbial Ecology*; 2016; 460–469. <https://doi.org/10.1007/s00248-016-0797-6> PMID: 27278778
11. Bletz MC, Bina Perl RG, Vences M. Skin microbiota differs drastically between co-occurring frogs and newts. *R Soc Open Sci*. 2017; 4. <https://doi.org/10.1098/rsos.170107> PMID: 28484639
12. Bletz MC, Archer H, Harris RN, McKenzie VJ, Rabemananjara FCE, Rakotoarison A, et al. Host ecology rather than host phylogeny drives amphibian skin microbial community structure in the biodiversity hotspot of Madagascar. *Front Microbiol*. 2017; 8: Article 1530. <https://doi.org/10.3389/fmicb.2017.01530> PMID: 28861051
13. Belden LK, Hughey MC, Rebollar EA, Umile TP, Loftus SC, Burzynski EA, et al. Panamanian frog species host unique skin bacterial communities. *Front Microbiol*. 2015; 6: 1171. <https://doi.org/10.3389/fmicb.2015.01171> PMID: 26579083
14. Abarca JG, Vargas G, Zuniga I, Whitfield SM, Woodhams DC, Kerby J, et al. Assessment of bacterial communities associated with the skin of costa rican amphibians at la selva biological station. *Front Microbiol*. 2018; 9: 1–12. <https://doi.org/10.3389/fmicb.2018.00001>
15. Bletz MCMC, Myers J, Woodhams DCDC, Rabemananjara FCEFCE, Rakotonirina A, Weldon C, et al. Estimating herd immunity to amphibian chytridiomycosis in Madagascar based on the defensive function of amphibian skin bacteria. *Front Microbiol*. 2017; 8. <https://doi.org/10.3389/fmicb.2017.01751> PMID: 28959244
16. Blümel S, Mark B, Busse H-J, Kämpfer P, Stolz A. *Pigmentiphaga kullae* gen. nov., sp. nov., a novel member of the family Alcaligenaceae with the ability to decolorize azo dyes aerobically. *Int J Syst Evol Microbiol. Microbiology Society*; 2001; 51: 1867–1871. <https://doi.org/10.1099/00207713-51-5-1867> PMID: 11594620
17. Yoon J-H, Kang S-J, Kim W, Oh T-K. *Pigmentiphaga daeguensis* sp. nov., isolated from wastewater of a dye works, and emended description of the genus *Pigmentiphaga*. *Int J Syst Evol Microbiol. Microbiology Society*; 2007; 57: 1188–1191. <https://doi.org/10.1099/ijls.0.64901-0> PMID: 17551027
18. Lee J-J, Srinivasan S, Kim MK. *Pigmentiphaga soli* sp. nov., a bacterium isolated from soil. *J Microbiol. Springer*; 2011; 49: 857–861. <https://doi.org/10.1007/s12275-011-1375-8> PMID: 22068507
19. Lee SD. *Pigmentiphaga aceris* sp. nov., isolated from tree sap. *Int J Syst Evol Microbiol. Microbiology Society*; 2017; 67: 3198–3202. <https://doi.org/10.1099/ijsem.0.002073> PMID: 28840799
20. Chen Y-G, Zhang Y-Q, Huang K, Tang S-K, Cao Y, Shi J-X, et al. *Pigmentiphaga litoralis* sp. nov., a facultatively anaerobic bacterium isolated from a tidal flat sediment. *Int J Syst Evol Microbiol. Microbiology Society*; 2009; 59: 521–525. <https://doi.org/10.1099/ijls.0.002949-0> PMID: 19244433
21. Proença DN, Grass G, Morais P V. Understanding pine wilt disease: roles of the pine endophytic bacteria and of the bacteria carried by the disease-causing pinewood nematode. *Microbiologyopen*. Wiley Online Library; 2017; 6: e00415.
22. Bridger N, Drews S, Burdz T, Wiebe D, Pacheco AL, Ng B, et al. Isolation and characterization of *Pigmentiphaga*-like isolates from human clinical material. *J Med Microbiol. Microbiology Society*; 2013; 62: 708–711. <https://doi.org/10.1099/jmm.0.051615-0>

23. Bernier A-M, Bernard K. Draft Whole-Genome Sequences for Two *Pigmentiphaga* Isolates Recovered from Human Clinical Materials. *Genome Announc. Am Soc Microbiol*; 2017; 5: e00726–17. <https://doi.org/10.1128/genomeA.00726-17> PMID: 28774981
24. Huang J, Ling J, Kuang C, Chen J, Xu Y, Li Y. Microbial biodegradation of aniline at low concentrations by *Pigmentiphaga daeguensis* isolated from textile dyeing sludge. *Int Biodeterior Biodegradation*. Elsevier; 2018; 129: 117–122.
25. Garrido-Sanz D, Manzano J, Martín M, Redondo-Nieto M, Rivilla R. Metagenomic analysis of a biphenyl-degrading soil bacterial consortium reveals the metabolic roles of specific populations. *Front Microbiol. Frontiers*; 2018; 9: 232.
26. Amir A, McDonald D, Navas-Molina JA, Kopylova E, Morton JT, Zech Xu Z, et al. Deblur rapidly resolves single-nucleotide community sequence patterns. Gilbert JA, editor. *mSystems*. 2017; 2.
27. Wang Q, Garrity GM, Tiedje JM, Cole JR. Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Appl Environ Microbiol*. 2007; 73: 5261–5267. <https://doi.org/10.1128/AEM.00062-07> PMID: 17586664
28. Caporaso JG, Lauber CL, Walters WA, Berg-lyons D, Lozupone CA, Turnbaugh PJ, et al. Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. *Proc Natl Acad Sci*. 2011; 108: 4516–4522. <https://doi.org/10.1073/pnas.1000080107> www.pnas.org/cgi/doi/10.1073/pnas.1000080107 PMID: 20534432
29. Hedges SB, Dudley J, Kumar S. TimeTree: a public knowledge-base of divergence times among organisms. *Bioinformatics*. 2006; 22: 2971–2972. Available: <https://doi.org/10.1093/bioinformatics/btl505> PMID: 17021158
30. Wickham H. *ggplot2: Elegant graphics for data analysis*. New York: Springer; 2009.
31. Li H, Durbin R. Fast and accurate long-read alignment with Burrows–Wheeler transform. *Bioinformatics*. Oxford University Press; 2010; 26: 589–595. <https://doi.org/10.1093/bioinformatics/btp698> PMID: 20080505
32. Koboldt DC, Zhang Q, Larson DE, Shen D, McLellan MD, Lin L, et al. VarScan 2: somatic mutation and copy number alteration discovery in cancer by exome sequencing. *Genome Res*. Cold Spring Harbor Lab; 2012; 22: 568–576. <https://doi.org/10.1101/gr.129684.111> PMID: 22300766
33. Seemann T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics*. Oxford University Press; 2014; 30: 2068–2069. <https://doi.org/10.1093/bioinformatics/btu153> PMID: 24642063
34. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, et al. NCBI prokaryotic genome annotation pipeline. *Nucleic Acids Res*. Oxford University Press; 2016; 44: 6614–6624. <https://doi.org/10.1093/nar/gkw569> PMID: 27342282
35. Darling ACE, Mau B, Blattner FR, Perna NT. Mauve: multiple alignment of conserved genomic sequence with rearrangements. *Genome Res*. Cold Spring Harbor Lab; 2004; 14: 1394–1403. <https://doi.org/10.1101/gr.2289704> PMID: 15231754
36. Kanehisa M, Sato Y, Morishima K. BlastKOALA and GhostKOALA: KEGG tools for functional characterization of genome and metagenome sequences. *J Mol Biol*. Elsevier; 2016; 428: 726–731. <https://doi.org/10.1016/j.jmb.2015.11.006> PMID: 26585406
37. Blin K, Wolf T, Chevrette MG, Lu X, Schwalen CJ, Kautsar SA, et al. antiSMASH 4.0—improvements in chemistry prediction and gene cluster boundary identification. *Nucleic Acids Res*. Oxford University Press; 2017; 45: W36–W41. <https://doi.org/10.1093/nar/gkx319> PMID: 28460038
38. Weber T, Rausch C, Lopez P, Hoof I, Gaykova V, Huson DH, et al. CLUSEAN: a computer-based framework for the automated analysis of bacterial secondary metabolite biosynthetic gene clusters. *J Biotechnol*. Elsevier; 2009; 140: 13–17. <https://doi.org/10.1016/j.jbiotec.2009.01.007> PMID: 19297688
39. de Jong A, van Heel AJ, Kok J, Kuipers OP. BAGEL2: mining for bacteriocins in genomic data. *Nucleic Acids Res*. Oxford University Press; 2010; 38: W647–W651. <https://doi.org/10.1093/nar/gkq365> PMID: 20462861
40. Starcevic A, Zucko J, Simunkovic J, Long PF, Cullum J, Hranueli D. ClustScan: an integrated program package for the semi-automatic annotation of modular biosynthetic gene clusters and in silico prediction of novel chemical structures. *Nucleic Acids Res*. Oxford University Press; 2008; 36: 6882–6892. <https://doi.org/10.1093/nar/gkn685> PMID: 18978015
41. Caboche S, Pupin M, Leclère V, Fontaine A, Jacques P, Kucherov G. NORINE: a database of nonribosomal peptides. *Nucleic Acids Res*. Oxford University Press; 2007; 36: D326–D331. <https://doi.org/10.1093/nar/gkm792> PMID: 17913739
42. Katoh K, Rozewicki J, Yamada KD. MAFFT online service: multiple sequence alignment, interactive sequence choice and visualization. *Brief Bioinform*. 2017;
43. Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. *Mol Biol Evol*. 2016; 33: msw054. <https://doi.org/10.1093/molbev/msw054> PMID: 27004904



44. Carver T, Thomson N, Bleasby A, Berriman M, Parkhill J. DNAPlotter: circular and linear interactive genome visualization. *Bioinformatics*. Oxford University Press; 2008; 25: 119–120. <https://doi.org/10.1093/bioinformatics/btn578> PMID: 18990721
45. Crosa JH, Walsh CT. Genetics and assembly line enzymology of siderophore biosynthesis in bacteria. *Microbiol Mol Biol Rev. Am Soc Microbiol*; 2002; 66: 223–249. <https://doi.org/10.1128/MMBR.66.2.223-249.2002> PMID: 12040125
46. Riley MA, Wertz JE. Bacteriocins: evolution, ecology, and application. *Annu Rev Microbiol. Annual Reviews* 4139 El Camino Way, PO Box 10139, Palo Alto, CA 94303–0139, USA; 2002; 56: 117–137. <https://doi.org/10.1146/annurev.micro.56.012302.161024> PMID: 12142491
47. Kumariya R, Garsa AK, Rajput YS, Sood SK, Akhtar N, Patel S. Bacteriocins: Classification, synthesis, mechanism of action and resistance development in food spoilage causing bacteria. *Microb Pathog*. Elsevier; 2019;
48. Cotter PD, Ross RP, Hill C. Bacteriocins—a viable alternative to antibiotics? *Nat Rev Microbiol*. Nature Publishing Group; 2013; 11: 95. <https://doi.org/10.1038/nrmicro2937> PMID: 23268227
49. Robinson SL, Christenson JK, Wackett LP. Biosynthesis and chemical diversity of  $\beta$ -lactone natural products. *Nat Prod Rep*. Royal Society of Chemistry; 2019;
50. Czech L, Höppner A, Kobus S, Seubert A, Riclea R, Dickschat JS, et al. Illuminating the catalytic core of ectoine synthase through structural and biochemical analysis. *Sci Rep*. Nature Publishing Group; 2019; 9: 364. <https://doi.org/10.1038/s41598-018-36247-w> PMID: 30674920
51. Schulz S, Dickschat JS. Bacterial volatiles: the smell of small organisms. *Nat Prod Rep*. Royal Society of Chemistry; 2007; 24: 814–842. <https://doi.org/10.1039/b507392h> PMID: 17653361
52. Zhang L, Khabbaz SE, Wang A, Li H, Abbasi PA. Detection and characterization of broad-spectrum antipathogen activity of novel rhizobacterial isolates and suppression of *Fusarium* crown and root rot disease of tomato. *J Appl Microbiol*. 2015; 118: 685–703. <https://doi.org/10.1111/jam.12728> PMID: 25512025
53. Ossowicki A, Jafra S, Garbeva P. The antimicrobial volatile power of the rhizospheric isolate *Pseudomonas donghuensis* P482. *PLoS One*. 2017; 12: 1–13. <https://doi.org/10.1371/journal.pone.0174362> PMID: 28358818